

## For healthcare providers treating children 18 years of age and younger

### HEALTHCARE PROVIDERS SHOULD:

#### ASSESS.

Conduct a physical examination to identify findings that:

- Suggest more severe TBI (e.g., hemotympanum, pupillary asymmetry).
- May impact management of mTBI (e.g., concurrent injuries or baseline deficits, oculomotor dysfunction).
- Suggest other contributions to symptoms (e.g., dehydration, cervical tenderness, scalp hematoma).

Do not image routinely (including CT & MRI).

- Use validated clinical decision rules predicting risk for more severe injury to determine need.

Assess symptoms using validated scales. Consider cognitive and balance testing.

Conduct a history to identify risk factors for poor prognosis using validated prediction rules.

#### COUNSEL.

Provide information about:

- Warning signs that injury may be more serious.
- Typical recovery course.
- How to prevent further injury.
- Gradual re-introduction of activity that does not worsen symptoms.
- The need for social and emotional support.

Offer clear instructions (preferably verbal and written) on return to activity, including school and sports, customized to the patient's symptoms.

- After a few days of rest (2-3 days), begin light activity & then gradually re-introduce regular activities (not inclusive of sports) that do not significantly worsen symptoms.
- Assess school-related needs & monitor progress in collaboration with parents and school professionals.
- Once back to regular non-sports activities (including school), patient can begin return to sports using a standard progression with gradually increasing levels of physical exertion.
- No return to contact sports activity until symptom-free with exertion (including without the use of pain medication).

#### REFER.

Identify and tailor treatment plans/referrals to address:

- Acutely worsening symptoms → consider neuroimaging.
- Chronic headache → non-opioid analgesia (monitor for overuse), multi-disciplinary evaluation.
- Vestibulo-ocular dysfunction → vestibular rehabilitation.
- Worsening sleep problem → sleep hygiene, sleep specialist.
- Cognitive impairment → treatment directed at etiology, neuropsychological evaluation.
- Emotional dysfunction → psychotherapeutic evaluation and treatment.

**A combination of risk factors that may indicate need for neuroimaging include:**

- Age < 2 years old
- Recurrent vomiting
- Loss of consciousness
- Severe mechanism of injury
- Severe or worsening headache
- Amnesia
- Non-frontal scalp hematoma
- Glasgow Coma Score < 15
- Clinical suspicion for skull fracture

**Examples of validated scales include, but aren't limited to:**

- Post-Concussion Symptom Scale
- Health and Behavior Inventory
- Post-Concussion Symptom Inventory
- Acute Concussion Evaluation

**Factors associated with poor prognosis:**

- Older age (older children/adolescents) or Hispanic ethnicity
- Lower socio-economic status
- History of intracranial injury
- Premorbid histories of mTBI or increased pre-injury symptoms
- Neurological or psychiatric disorder
- Learning difficulties or lower cognitive ability
- Family and social stressors

**Parents should watch for warning signs:**

- A headache that gets worse & does not go away
- Significant nausea or repeated vomiting
- Increased confusion, restlessness, or agitation
- Slurred speech, drowsiness, or inability to wake up
- Weakness, numbness, or decreased coordination
- Loss of consciousness, convulsions, or seizures

**Steps in a return to play progression generally include:**

- Step 1: Return to regular non-sports activities
- Step 2: Light aerobic exercise
- Step 3: Sport-specific exercise
- Step 4: Non-contact training drills
- Step 5: Full contact practice
- Step 6: Return to sport

**Refer patients whose symptoms do not resolve as expected with standard care after 4-6 weeks.**

