

PrEP choice	Formulation	Dosing Frequency	Gender & Weight	CrCl	Baseline Labs	Every 3mo	Every 6mo	Every 12mo
Oral Truvada	<ul style="list-style-type: none"> 200mg Emtricitabine-300mg Tenofovir disoproxil Generic available 	<ul style="list-style-type: none"> Once daily 2-1-1 for MSM only 	<ul style="list-style-type: none"> All genders Weight ≥ 35kg 	<ul style="list-style-type: none"> ≥ 60 	<ul style="list-style-type: none"> 4th gen Ab/Ag HIV test STI testing based on what sites are being used for sex Hep B (HBsAg, HBsAb & HBcAb) Hep C Ab if MSM, TGW or PWID CrCl Pregnancy test for Truvada Lipid panel for Descovy 	<ul style="list-style-type: none"> 4th gen Ab/Ag and HIV RNA PCR testing STI testing for MSM & TGW 	<ul style="list-style-type: none"> 4th gen Ab/Ag and HIV RNA PCR testing STI testing for everyone based on sites being used for sex CrCL if baseline was <90 	<ul style="list-style-type: none"> CrCL if baseline was >90 Hep C for MSM, TGW and PWID Lipid panel for Descovy
Oral Descovy	<ul style="list-style-type: none"> 200mg Emtricitabine-25mg Tenofovir alafenamide Generic coming soon 	<ul style="list-style-type: none"> Once daily 	<ul style="list-style-type: none"> NOT approved for receptive vaginal sex Weight ≥ 35kg 	<ul style="list-style-type: none"> ≥ 30 	<ul style="list-style-type: none"> 4th gen Ab/Ag HIV test STI testing based on what sites are being used for sex Hep B (HBsAg, HBsAb & HBcAb) Hep C Ab if MSM, TGW or PWID CrCl Pregnancy test for Truvada Lipid panel for Descovy 	<ul style="list-style-type: none"> 4th gen Ab/Ag and HIV RNA PCR testing STI testing for MSM & TGW 	<ul style="list-style-type: none"> 4th gen Ab/Ag and HIV RNA PCR testing STI testing for everyone based on sites being used for sex CrCL if baseline was <90 	<ul style="list-style-type: none"> CrCL if baseline was >90 Hep C for MSM, TGW and PWID Lipid panel for Descovy
Injectable Apretude	<ul style="list-style-type: none"> Single-dose intramuscular injection of 200mg/mL of Cabotegravir 	<ul style="list-style-type: none"> Once every 2 months 	<ul style="list-style-type: none"> All genders Adult and adolescents Weighing ≥ 35kg 		<ul style="list-style-type: none"> Get an HIV-1 viral load (HIV-1 RNA Quantitative PCR) then refer to SIS or other Apretude provider 	<ul style="list-style-type: none"> SIS or other Apretude provider will repeat monitoring at 3, 6, and 12 months 		
Injectable Yeztugo	<ul style="list-style-type: none"> 927mg by subcutaneous injection of Lenacapavir (2 x 1.5 mL injections) 	<ul style="list-style-type: none"> Once every 6 months 	<ul style="list-style-type: none"> All genders Adult and adolescents Weighing ≥ 35kg 		<ul style="list-style-type: none"> Get an HIV-1 viral load (HIV-1 RNA Quantitative PCR) then refer to SIS or other Yeztugo provider 	<ul style="list-style-type: none"> SIS or other Yeztugo provider will repeat monitoring at 3, 6, and 12 months 		

MSM: Men who have sex with men, TGW: Transgender women, PWID: Persons who inject drugs, LAD: Lymphadenopathy